

# Translational Animal Models for Liver Cancer

Michele Obeid<sup>1</sup>, Ramzy C. Khabbaz<sup>1</sup>, Kelly D. Garcia<sup>2</sup>, Kyle M. Schachtschneider<sup>3</sup>,  
Ron C. Gaba<sup>1</sup>

<sup>1</sup>Department of Radiology, University of Illinois, 1740 West Taylor Street MC 931, Chicago, IL, 60612, United States, <sup>2</sup>College of Medicine, University of Illinois, 1740 West Taylor Street MC 931, Chicago, IL, 60612, United States, <sup>3</sup>Department of Biological Resources Laboratory, University of Illinois, 1740 West Taylor Street MC 931, Chicago, IL, 60612, United States



## Corresponding Author:

Ron Gaba, Department of Radiology, University of Illinois, 1740 West Taylor Street MC 931, Chicago, IL, 60612, United States.  
E-mail: rongaba@yahoo.com

Received: 12-December-2017

Accepted: 10-February-2018

Published: 24-February-2018

## ABSTRACT

Animal models have become increasingly important in the study of hepatocellular carcinoma (HCC), as they serve as a critical bridge between laboratory-based discoveries and human clinical trials. Developing an ideal animal model for translational use is challenging, as the perfect model must be able to reproduce human disease genetically, anatomically, physiologically, and pathologically. This brief review provides an overview of the animal models currently available for translational liver cancer research, including rodent, rabbit, non-human primate, and pig models, with a focus on their respective benefits and shortcomings. While small animal models offer a solid starting point for investigation, large animal HCC models are becoming increasingly important for translation of preclinical results to clinical practice.

**Key words:** Animal models, hepatocellular carcinoma, translational research

## INTRODUCTION

**H**epatocellular carcinoma (HCC), or primary liver cancer, is the fifth most commonly diagnosed cancer and the second most common cause of cancer death in men worldwide, while it is the seventh most commonly diagnosed cancer and the sixth most common cause of cancer death in women.<sup>[1,2]</sup> To better understand the natural history and treatment of this deadly disease, animal models have become increasingly important, as they serve as a critical bridge between laboratory-based discoveries and human clinical trials. Historically, animal liver cancer models have spanned rodents,<sup>[3]</sup> rabbits,<sup>[4]</sup> dogs,<sup>[5]</sup> monkeys,<sup>[6]</sup> and pigs.<sup>[7]</sup> Over the past several decades, however, existing

animal models of liver cancer have been refined to more accurately represent the disease, and new animal models of HCC have been developed. Mammalian models currently utilized for investigation of HCC include mouse, rat, woodchuck, rabbit, and porcine platforms,<sup>[8]</sup> all employed with the intent of allowing the accurate preclinical study of disease development, detection, treatment, and progression.<sup>[8]</sup> This brief review provides an overview of the animal models currently available for translational liver cancer research, with a focus on their respective benefits and shortcomings.

### The ideal animal model

Developing a robust animal model that is suitable for translational use is challenging, as the ideal model must be able to reproduce human disease genetically, anatomically, physiologically, and pathologically.<sup>[9]</sup> A perfect animal model should harbor disease that arises from a relevant cell line which lends itself to propagation, characterization, storage, and study *in vitro*, which allows for benchtop molecular assays to be performed before or in parallel to animal experiments. It should mimic human disease on the molecular, cellular,

### Access this article online

Quick Response Code:



Website:  
www.americanjir.com

DOI:  
10.25259/AJIR-11-2017

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

and phenotypic basis, be reliable and predictable in tumor generation, proliferation, and growth kinetics, develop in a practical time frame, be readily imaged, allow for accurate assessment of treatment effects (i.e., no spontaneous tumor necrosis), and manifest survival differences in affected versus non-affected subjects. Furthermore, preclinical tumors should be generated in a background setting that replicates the clinical context in which the disease is found in humans (e.g., liver cirrhosis for HCC) to optimally recapitulate the tumor-host microenvironment on a molecular basis and to best reflect tumor and local host organ interactions. To this end, autochthonous tumor models those that spontaneously arise in the organ of interest rather than being transplanted (orthotopic model) confer potential advantages in studying the molecular basis of disease. Suitable animal models should be of adequate size to allow for utilization of tools employed in clinical practice. Finally, an ideal animal model should not be cost prohibitive or logistically impractical and should minimize pain and suffering experienced by the animals under study.

## Rodent models

### Mouse models

There are several methods for induction of murine liver cancer. These include chemical induction (through N-nitrosodiethylamine or diethylnitrosamine [DEN], aflatoxin B1, carbon tetrachloride, choline deficiency, or thioacetamide exposure), xenografting (e.g., human tumor cell lines or fragments implanted into immunodeficient animals), and genetic modification (e.g., hepatitis B and C virus transgenic mice expressing specific portions of the viral genome).<sup>[3]</sup>


Small animals like mice offer advantages for cancer studies (Table 1). These include ease of handling, low procurement and maintenance costs, established methods for biologic and genetic manipulation, and the potential for high throughput. Two additional major advantages include rapid tumor induction, as mouse cell lines have very high rates of tumor growth, and easy surveillance of tumor growth by direct visualization, palpation, and measurement (Figure 1). For example, in the mouse xenograft model, the time span between tumor cell injection and tumor development is relatively short (days to weeks),<sup>[10]</sup> which although efficient, does not mimic human HCC growth kinetics.<sup>[11]</sup>

Even though mice provide versatile and flexible animal models, they have several limitations (Table 1).<sup>[12]</sup> First, researchers need to consider that cellular origin and signaling pathways differ between mice and humans. For example, murine strains are highly inbred, essentially having 100% homozygosity at every locus.<sup>[13,14]</sup> Murine cells also require fewer genetic mutations to develop tumorigenic phenotypes, making them easier to immortalize than human cells.<sup>[15]</sup> In addition, unlike human cells, their signaling pathway requirements are reduced in Ras oncogene transformation,<sup>[16]</sup> and in contrast to humans,

**Table 1:** Rodent tumor model advantages and disadvantages

Advantages	Drawbacks
Ease of handling	Genetic differences
Low procurement and maintenance costs	Small size precludes clinical device utilization and selective intra-arterial therapy
High throughput	Difficult to induce native HCC
Rapid tumor induction	Questionable translatability to human disease
Easy surveillance of tumor growth	
Established methods for biologic and genetic manipulation	

HCC: Hepatocellular carcinoma



**Figure 1:** Photographs demonstrate mouse subcutaneous hepatocellular carcinoma xenograft (arrow) before (a) and following (b) euthanasia and tumor harvest.

*Mus musculus* demonstrate high levels of telomerase activity in normal somatic tissues.<sup>[15,17]</sup> Another major drawback of employing mouse models is their small size, which precludes clinical device utilization and limits technical capability for interventions such as vascular catheterization for the study of transarterial therapy. Furthermore, without genetic modification, mice tend to develop sarcomas and lymphomas, whereas humans have a bias toward developing epithelial cancers, or carcinomas, with age.<sup>[15]</sup> Specific to HCC research, mice are not susceptible to hepatitis viral infection. In addition, though mice develop liver steatosis when exposed to a diet high in fats or alcohol, the steatosis does not progress to a translational non-alcoholic steatohepatitis, cirrhosis, or HCC model. Finally, recent literature has uncovered meaningful issues with reproducibility of findings made in murine systems,<sup>[18]</sup> and the direct translatability of findings made in mouse models to human clinical practice is questionable. To this end, while murine models are commonly used in preclinical drug research, many agents that show promise in mouse studies ultimately fail in human clinical trials.<sup>[19]</sup>


### Rat models

Among rat HCC models, the most frequently employed are the MCA-RH 7777 and N1-S1 cell lines derived from the Morris<sup>[20]</sup> and Novikoff<sup>[21]</sup> hepatoma models, respectively. The Morris hepatoma model was generated in Buffalo

rats following exposures to N-2-fluorenylphthalamic acid, which yielded a tumor used to develop the MCA-RH 7777 cell line.<sup>[20]</sup> Tumor development using this line consists of intra-hepatic injection of syngeneic tumor cells in Buffalo rats, although the model is compatible with Sprague-Dawley rats as well.<sup>[22]</sup> Advantages of this model include high inoculation rate and  $\alpha$ -fetoprotein production, which allows for therapeutic monitoring. Shortcomings include lack of Buffalo rat availability, variability on imaging and histological exam,<sup>[23]</sup> and aggressive metastatic behavior (although this may be useful in modeling multifocal HCC).<sup>[24]</sup> The Novikoff hepatoma model originated from exposure of Sprague-Dawley rats to 4-dimethylaminoazobenzene yielding the N1-S1 cell line (isolated from ascitic fluid). N1-S1 tumors are also produced through intra-hepatic injection of syngeneic tumor cells. The advantage of this model includes the ability to generate concurrent liver cirrhosis through common bile duct ligation,<sup>[25]</sup> while it is limited by suboptimal tumor induction rates,<sup>[26]</sup> and potential for spontaneous regression.<sup>[22]</sup>

Recently, a DEN-induced Wistar rat autochthonous HCC model was developed (Figure 2).<sup>[27]</sup> Its benefits include recapitulation of the hepatocyte injury cirrhosis malignancy evolutionary cycle as is seen with human HCC, the presence of liver disease (which theoretically renders subjects susceptible to known complications associated with hepatic insufficiency), and tumor hypervascularity. In addition, the presence of cirrhosis and hepatic dysfunction narrows the therapeutic window for therapy compared to models without background liver disease,<sup>[27]</sup> as is seen in the human condition. The main limitation of this model is the 3-month induction time required for proper tumor development.<sup>[27]</sup>

In considering procedure based therapeutic interventions, transarterial treatments may be performed in rat HCC models,



**Figure 2:** Photograph reveals multifocal disease (arrows) in diethylnitrosamine-induced Wistar rat autochthonous hepatocellular carcinoma model (image courtesy of Terence P. Gade M.D. Ph.D., Department of Radiology, University of Pennsylvania).

but most require a laparotomy and do not entail selective therapy. A segmental approach to transarterial therapy was described in the DEN-induced Wistar rat autochthonous HCC model but was associated with lengthy procedure times in undertaking selective arteriography.<sup>[27]</sup>

### Woodchuck models

From 1950 to 1970, the Woodchuck was introduced as a viral hepatitis model.<sup>[28]</sup> Unlike human hepatitis virus, the inoculated Woodchuck hepatitis virus (WHV) does not incite cirrhosis in woodchucks.<sup>[8]</sup> In addition, these animals are more challenging to handle and do not breed as efficiently as other laboratory rodents. However, WHV does result in spontaneous generation of HCC, allowing for utilization of this model for studying transarterial and percutaneous ablative therapy of primary HCC tumors in pre-clinical settings.<sup>[29]</sup> More recent studies have demonstrated a similarity between woodchuck and human liver arterial anatomy, and that femoral arterial access can be achieved in the woodchuck for catheterization of the hepatic arterial system (as in human clinical practice), although this procedure is somewhat challenging due to the small caliber of the common femoral artery.<sup>[30]</sup> The major drawback of this animal model is the prolonged time period (1–4 years) required to develop liver tumors of 1000 cm<sup>3</sup>,<sup>[29]</sup> which is impractical within the desired time frame of scientific investigation.

### Rabbit models

Developed in the 1930s by Rous and Beard,<sup>[31]</sup> the VX2 tumor is an anaplastic squamous cell carcinoma which is induced in rabbits. Tumor development takes only 2–3 weeks, which makes the VX2 model an efficient model for the study of liver tumors. This model offers several advantages (Table 2), including reliability of tumor induction by orthotopic allografting, rapid growth, easy propagation in skeletal muscle, and ability to allograft donor tumors into multiple recipient rabbits.<sup>[32]</sup> Many interventions can be accomplished utilizing the VX2 platform, including both transarterial and ablative therapies.<sup>[33]</sup> Liver tumors grown from VX2 cell lines are well suited for investigating ablative therapies given sonographic visibility, and the innate hypervascularity of VX2 tumors coupled with the relatively more sizeable rabbit vasculature permits technically easier investigation of transarterial locoregional treatments for liver tumors (Figure 3). *In vitro* growth of VX2 cell lines allow cytotoxicity investigations, which may be correlated to results obtained *in vivo*.<sup>[34]</sup> Limitations of the VX2 model (Table 2) includes squamous cell origin (dissimilar to HCC, which is an adenocarcinoma), unknown tumor biology, peripheral vascularization, varying tumor kinetics, and unknown genome organization. Another critical shortcoming of this model is spontaneous central tumor necrosis, which confounds the evaluation of therapeutic efficacy after pharmacological or interventional treatment. Since its introduction, progress has been made to



**Figure 3:** Fluoroscopic spot image from hepatic arteriogram during locoregional therapy of rabbit VX2 liver tumor. Left hepatic arteriogram demonstrates hypervascular left hepatic lobe tumor (arrow).

increase the viability of VX2 tumors by creating modified lines,<sup>[35]</sup> potentially increasing the platform's general utility. The dissimilarity of VX2 tumor biology from human HCC also precludes definitive investigation of cytotoxic agents or targeted therapeutics, which represents a major drawback in this era of precision medicine.

### Companion animal and non-human primate models

Canine cancer models have been used to study osteosarcoma, sarcoma, lung cancer, nasal and oral malignancy, and malignant non-Hodgkin's lymphoma given a high analogy to the corresponding human tumors.<sup>[36]</sup> However, dogs tend to develop more lymphoid and sarcoma tumor types rather than carcinomas, and compared to humans, they have varying drug sensitivity. In addition, dogs are patients, not experimental animals, and therefore, cannot be systematically studied in the same way other animals can be. Thus, they are not widely used in the preclinical study of liver cancer. Non-human primates have genetic, anatomical, and physiological similarity to humans, offering advantages for studying cancer,<sup>[37]</sup> rhesus macaques, cynomolgus macaques, and common marmosets have been shown to be susceptible to HCC induction.<sup>[38]</sup> However, these animals have relatively small size, and metabolic studies have shown that non-human primates do not always simulate human cancer or disease.<sup>[8]</sup> Pertaining to these animals, the ethical concerns of using companion animals and non-human primates for translational research could substantially limit the preclinical advancement of HCC models in these systems.

### Pig models

Swine cancer models have shown to be a promising animal model (Table 3) due to their similarities to humans in size,

**Table 2:** Rabbit VX2 tumor model advantages and disadvantages

Advantages	Drawbacks
Reliable tumor induction by orthotopic allograft implantation	Physiologically different host
Rapid growth	Genetically dissimilar
Easy propagation in skeletal muscle	Varying tumor kinetics
Ability to graft donor tumors into multiple recipient rabbits	Spontaneous necrosis
Innate hypervascularity	Non-diseased liver
More sizeable vasculature than rodent models	

**Table 3:** Porcine model advantages and disadvantages

Advantages	Disadvantages
Similar size and anatomy allows utilization of same devices and instruments employed in clinical practice, enabling translation to human clinical trials	Larger housing requirements compared to small animals
Similar physiology allows for consistent pharmacokinetic analysis of drug metabolism	Longer generation intervals
Similar genetic background allows relevant investigation of molecular basis of disease and drug therapy	Lack of pig model of hepatitis infection
	Lower quality genome and fewer genomic tools compared to mice and humans

anatomy, pathophysiology, metabolism, genetics, epigenetics, and pathology.<sup>[39]</sup> They are less expensive than non-human primate models, and they age 3–5 times faster than humans.<sup>[40]</sup> This life cycle permits enough time to develop, characterize, and modulate cancer in a pig model.<sup>[41]</sup>


For years, swine have served as a valuable resource for procedural training in surgery and interventional radiology. The size and anatomy of the pig liver, which is similar to that of humans, allows for utilization of instruments and devices employed in the care of patients, enabling easier application and translation to human clinical trials.<sup>[42]</sup> Unlike other animal models, pigs also exhibit similar genetics and drug metabolism to humans,<sup>[43]</sup> which allows for relevant investigation of the molecular basis of disease in addition to drug pharmacokinetics and therapeutic analyses. Until recently, pigs were most commonly used for practicing techniques rather than modeling disease; however, the establishment of porcine HCC models shows great promise in advancing diagnostic and therapeutic discoveries for liver cancer.

Drug-induced pig models have been created in which DEN was introduced intraperitoneally, weekly, for a period of 3–4 months.<sup>[44,45]</sup> These models developed cirrhosis and, eventually, liver cancer over a 12–24 months period; however, this time frame is relatively impractical given the expense of housing animals for a protracted period, the unpredictability of spontaneous cancer formation, and the resultant delay in initiation of tumor-related scientific activities. In addition, tumors formed in the described model were small and numerous, which is more representative of advanced human liver cancer where the disease has already disseminated.<sup>[8]</sup>

The recently developed oncopig cancer model (OCM) is a unique transgenic pig model that develops the inducible site and cell-specific tumors after Cre recombinase exposure.<sup>[46]</sup> The OCM was designed to innately harbor mutations found in more than 50% of human cancers, *KRAS*<sup>G12D</sup> and *TP53*<sup>R167H</sup>, and the capability for rapid tumor induction in this model permits HCC study within a practical time frame.<sup>[46]</sup> In the OCM, the innate *KRAS*<sup>G12D</sup> and *TP53*<sup>R167H</sup> germline mutations are heterozygous in nature, closely modeling human disease.<sup>[46]</sup> OCM cohorts are developed through crossbreeding a transgene/major histocompatibility complex homozygous oncopig minnesota-mini composite sire with any number of domestic, mini-pig, or transgenic dams, allowing for the development of tumor-bearing pigs with a range of genetic backgrounds. This, coupled with the use of a maintenance diet, ensures manageable and clinically relevant animal subject sizes within study time periods. In initial investigations of the model, OCM HCC recapitulated human HCC histologic features, as well as transcriptional hallmarks of human HCC, including *TERT* reactivation, apoptosis evasion, angiogenesis activation, altered cell cycle regulation, and WNT signaling activation.<sup>[47]</sup> Autologous injection of porcine HCC cells into the OCM yielded tumors histologically characterized as HCC.<sup>[47]</sup> Importantly, liver cirrhosis can also be induced in swine models,<sup>[48]</sup> allowing evaluation of HCC in its native host environment. Because medical comorbidities related to the cancer-host environment may impact therapeutic capability and serve as a competing cause of mortality as in the case of liver cancer and hepatic cirrhosis the ability to induce comorbidities in the oncopig (Figure 4)<sup>[47]</sup> provides a unique opportunity to initiate tumors with and without comorbid illness to define combination-therapeutic approaches in a more controlled way that can be done in human patients. Finally, the OCM is immunocompetent, lending itself to the investigation of immunotherapies.<sup>[42]</sup> Given these attributes, the OCM has potential to serve as a valuable transitional bridge between preclinical small animal murine studies and human clinical trials.

## CONCLUSIONS

Animal models play a crucial role in serving as a bridge from preclinical HCC research to clinical care. Systematic



**Figure 4:** Trichrome stained oncopig liver histologic section after cirrhosis induction shows irregular hepatic lobules circumferentially surrounded by thick fibrous septa (arrows).

investigations in animal subjects may offer fundamental insights into mechanisms of disease development, as well as prevention, detection, treatment, and follow-up that can be applied to improve patient outcomes. While small animal models offer a solid starting point for investigation, the future will hold increasing importance for large animal HCC models as they become more widely used.

## Declaration of patient consent

Not required as there are no patients in this study.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

1. Stratton MR, Campbell PJ, Futreal PA. The cancer genome. *Nature* 2009;458:719-24.
2. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D, *et al.* Global cancer statistics. *CA Cancer J Clin* 2011;61:69-90.
3. Heindryckx F, Colle I, Van Vlierberghe H. Experimental mouse models for hepatocellular carcinoma research. *Int J Exp Pathol* 2009;90:367-86.
4. Reznik GK, Padberg G. Diethylnitrosamine-induced metastasizing hepatocellular carcinomas in New Zealand white rabbits. A tumor model for clinical investigations. *J Cancer Res Clin Oncol* 1991;117:123-9.

5. Hirao K, Matsumura K, Imagawa A, Enomoto Y, Hosogi Y. Primary neoplasms in dog liver induced by diethylnitrosamine. *Cancer Res* 1974;34:1870-82.
6. Lapis K, Sarosi I, Bocsi J, Thorgeirsson UP. Cytokeratin patterns of liver carcinomas induced by diethylnitrosamine in monkeys. *Lab Invest* 1995;72:748-59.
7. Flisikowska T, Kind A, Schnieke A. The new pig on the block: Modelling cancer in pigs. *Transgenic Res* 2013;22:673-80.
8. Aravalli RN, Steer CJ. Animal models of liver cancer. In: Aravalli RN, Steer CJ, editors. *Hepatocellular Carcinoma: Cellular and Molecular Mechanisms and Novel Therapeutic Strategies*. Cham: Springer International Publishing; 2014. p. 47-50.
9. Aravalli RN, Golzarian J, Cressman EN. Animal models of cancer in interventional radiology. *Eur Radiol* 2009;19:1049-53.
10. Rygaard J, Povsen CO. Heterotransplantation of a human malignant tumour to "nude" mice 1969. *APMIS* 2007;115:604-6.
11. Shao DM, Wang QH, Chen C, Shen ZH, Yao M, Zhou XD, *et al.* N-acetylglucosaminyltransferase V activity in metastatic models of human hepatocellular carcinoma in nude mice. *J Exp Clin Cancer Res* 1999;18:331-5.
12. Caviglia JM, Schwabe RF. Mouse models of liver cancer. *Methods Mol Biol* 2015;1267:165-83.
13. Holliday R. Neoplastic transformation: The contrasting stability of human and mouse cells. *Cancer Surv* 1996;28:103-15.
14. Kaiser J. The cancer test. *Science* 2015;348:1411-3.
15. Rangarajan A, Weinberg RA. Opinion: Comparative biology of mouse versus human cells: Modelling human cancer in mice. *Nat Rev Cancer* 2003;3:952-9.
16. Hamad NM, Elconin JH, Karnoub AE, Bai W, Rich JN, Abraham RT, *et al.* Distinct requirements for ras oncogenesis in human versus mouse cells. *Genes Dev* 2002;16:2045-57.
17. Kim SH, Kaminker P, Campisi J. Telomeres, aging and cancer: In search of a happy ending. *Oncogene* 2002;21:503-11.
18. Troublesome variability in mouse studies. *Nat Neurosci* 2009;12:1075.
19. Gould SE, Junttila MR, de Sauvage FJ. Translational value of mouse models in oncology drug development. *Nat Med* 2015;21:431-9.
20. Morris HP. Studies on the development, biochemistry, and biology of experimental hepatomas. *Adv Cancer Res* 1965;9:227-302.
21. Novikoff AB. A transplantable rat liver tumor induced by 4-dimethylaminoazobenzene. *Cancer Res* 1957;17:1010-27.
22. Cho HR, Choi JW, Kim HC, Song YS, Kim GM, Son KR, *et al.* Sprague-dawley rats bearing mcA-RH7777 cells for study of hepatoma and transarterial chemoembolization. *Anticancer Res* 2013;33:223-30.
23. Choi JW, Kim JH, Kim HC, Choi WS, Baek SY, Lee K, *et al.* Comparison of tumor vascularity and hemodynamics in three rat hepatoma models. *Abdom Radiol (NY)* 2016;41:257-64.
24. Guo Y, Klein R, Omary RA, Yang GY, Larson AC. Highly malignant intra-hepatic metastatic hepatocellular carcinoma in rats. *Am J Transl Res* 2010;3:114-20.
25. Thompson SM, Callstrom MR, Knudsen B, Anderson JL, Carter RE, Grande JP, *et al.* Development and preliminary testing of a translational model of hepatocellular carcinoma for MR imaging and interventional oncologic investigations. *J Vasc Interv Radiol* 2012;23:385-95.
26. Garin E, Denizot B, Roux J, Noiret N, Lepareur N, Moreau M, *et al.* Description and technical pitfalls of a hepatoma model and of intra-arterial injection of radiolabelled lipiodol in the rat. *Lab Anim* 2005;39:314-20.
27. Gade TP, Hunt SJ, Harrison N, Nadolski GJ, Weber C, Pickup S, *et al.* Segmental Transarterial Embolization in a Translational Rat Model of Hepatocellular Carcinoma. *J Vasc Interv Radiol* 2015;26:1229-37.
28. Tennant BC, Toshkov IA, Peek SF, Jacob JR, Menne S, Hornbuckle WE, *et al.* Hepatocellular carcinoma in the woodchuck model of hepatitis B virus infection. *Gastroenterology* 2004;127:S283-93.
29. Burke CT, Cullen JM, State A, Gadi S, Wilber K, Rosenthal M, *et al.* Development of an animal model for radiofrequency ablation of primary, virally induced hepatocellular carcinoma in the woodchuck. *J Vasc Interv Radiol* 2011;22:1613-80.
30. Wilkins LR, Stone JR, Mata J, Hawrylack A, Kubicka E, Brautigan DL. The use of the woodchuck as an animal model for evaluation of transarterial embolization. *J Vasc Interv Radiol* 2017;28:1467-71.
31. Rous P, Beard JW. The progression to carcinoma of virus-induced rabbit papillomas (Shope). *J Exp Med* 1935;62:523-48.
32. Galasko CS, Muckle DS. Intrasarcolemmal proliferation of the VX2 carcinoma. *Br J Cancer* 1974;29:59-65.
33. Parvinian A, Casadaban LC, Gaba RC. Development, growth, propagation, and angiographic utilization of the rabbit VX2 model of liver cancer: A pictorial primer and "how to" guide. *Diagn Interv Radiol* 2014;20:335-40.
34. Pascale F, Bedouet L, Baylatry M, Namur J, Laurent A. Comparative chemosensitivity of VX2 and HCC cell lines to drugs used in TACE. *Anticancer Res* 2015;35:6497-503.
35. Pascale F, Ghegediban SH, Bonneau M, Bedouet L, Namur J, Verret V, *et al.* Modified model of VX2 tumor overexpressing vascular endothelial growth factor. *J Vasc Interv Radiol* 2012;23:809-17.
36. Gardner HL, Fenger JM, London CA. Dogs as a model for cancer. *Annu Rev Anim Biosci* 2016;4:199-222.
37. Rogers J, Gibbs RA. Comparative primate genomics: Emerging patterns of genome content and dynamics. *Nat Rev Genet* 2014;15:347-59.
38. Foster JR. Spontaneous and drug-induced hepatic

- pathology of the laboratory beagle dog, the cynomolgus macaque and the marmoset. *Toxicol Pathol* 2005;33:63-74.
39. Swindle MM, Makin A, Herron AJ, Clubb FJ Jr., Frazier KS. Swine as models in biomedical research and toxicology testing. *Vet Pathol* 2012;49:344-56.
  40. Watson AL, Carlson DF, Largaespada DA, Hackett PB, Fahrenkrug SC. Engineered swine models of cancer. *Front Genet* 2016;7:78.
  41. Yeom SC, Cho SY, Park CG, Lee WJ. Analysis of reference interval and age-related changes in serum biochemistry and hematology in the specific pathogen free miniature pig. *Lab Anim Res* 2012;28:245-53.
  42. Schachtschneider KM, Schwind RM, Newson J, Kinachtchouk N, Rizko M, Mendoza-Elias N, *et al.* The oncopig cancer model: An innovative large animal translational oncology platform. *Front Oncol* 2017;7:190.
  43. Pollock CB, Rogatcheva MB, Schook LB. Comparative genomics of xenobiotic metabolism: A porcine-human PXR gene comparison. *Mamm Genome* 2007;18:210-9.
  44. Li X, Zhou X, Guan Y, Wang YX, Scutt D, Gong QY, *et al.* N-nitrosodiethylamine-induced pig liver hepatocellular carcinoma model: Radiological and histopathological studies. *Cardiovasc Intervent Radiol* 2006;29:420-8.
  45. Mitchell J, Tinkey PT, Avritscher R, Van Pelt C, Eskandari G, Konnath George S, *et al.* Validation of a preclinical model of diethylnitrosamine-induced hepatic neoplasia in yucatan miniature pigs. *Oncology* 2016;91:90-100.
  46. Schook LB, Collares TV, Hu W, Liang Y, Rodrigues FM, Rund LA, *et al.* A genetic porcine model of cancer. *PLoS One* 2015;10:e0128864.
  47. Schachtschneider KM, Schwind RM, Darfour-Oduro KA, De AK, Rund LA, Singh K, *et al.* A validated, transitional and translational porcine model of hepatocellular carcinoma. *Oncotarget* 2017;8:63620-34.
  48. Avritscher R, Wright KC, Javadi S, Uthamanthil R, Gupta S, Gagea M, *et al.* Development of a large animal model of cirrhosis and portal hypertension using hepatic transarterial embolization: A study in swine. *J Vasc Interv Radiol* 2011;22:1329-34.

**How to cite this article:** Obeid M, Khabbaz R, Garcia K, Schachtschneider K, Gaba R. Translational Animal Models for Liver Cancer. *Am J Interv Radiol* 2018, 2(2) 1-7.

AJIR