

Revascularization of Chronic Occlusive Disease of a Right Pulmonary Artery Graft

Russell Salamo¹, Nicholas Piening¹, Joseph Garrity MD¹, Ismael Moreno MD², Kirubahara Vaheesan MD¹

¹Department of Vascular and Interventional Radiology, Saint Louis University School of Medicine, 1402 S Grand Blvd, Saint Louis, MO, 63104, United States, ²Department of Vascular and Interventional Radiology, UC Davis, 4860 Y Street, Suite 3100, Sacramento, CA, 95817, United States



Corresponding Author:

Nicholas Piening, Department of Vascular and Interventional Radiology, Saint Louis University Medical School, 1402 S Grand Blvd, Saint Louis, MO, 63104, United States.
E-mail: npiening@slu.edu

Received: 30 March 2018

Accepted: 23 April 2018

Published: 27 April 2018

ABSTRACT

Anomalous aortic origin of the left coronary artery (AAOLCA) is a rare occurrence. This anomaly may lead to myocardial ischemia and, therefore, surgical repair is recommended to improve long-term survival in these patients. We present a case of successful mobilization of an anomalous left main coronary artery, pulmonary artery translocation, and interposition graft to the right pulmonary artery with a late complication of the right pulmonary artery graft occlusion. This case illustrates the difficulty in the recanalization of pulmonary artery graft occlusion and stenosis, and highlights the utility of treatment via a staged approach.

Keywords: Graft, Pulmonary artery, Revascularization, Thoracic, Thrombosis

INTRODUCTION

Anomalous aortic origin of a left coronary artery occurs when the left coronary artery (AAOLCA) arises from the incorrect coronary sinus. AAOLCA is a rare occurrence and may lead to myocardial ischemia and subsequent angina, dyspnea, and decreased exercise tolerance.^[1,2] Treatment of AAOLCA is surgical mobilization of an anomalous left main coronary artery, pulmonary artery

translocation, and interposition graft to the right pulmonary artery. With placement of stents, there is risk of fracture, risk of embolization, risk of stenosis with intra-stent neoproliferation, and the increased risk and complexity for further surgery.^[3] There remains limited data available regarding the incidence and management of pulmonary artery graft occlusion and stenosis.

CASE REPORT

A 41-year-old female presented with chest pain and shortness of breath. Her past medical history was significant for anomalous aortic origin of the left coronary artery (AAOLCA) status post-mobilization of the anomalous left main coronary artery, pulmonary artery translocation, and interposition of a 16 mm Dacron graft to the right pulmonary artery. Computed

Access this article online	
Quick Response Code: 	Website: www.americanjir.com
	DOI: 10.25259/AJIR-20-2018

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

tomography (CT) angiography of the chest showed an occlusion of the right pulmonary artery (Figure 1a). The patient was anticoagulated with aspirin and heparin, and interventional radiology was consulted for thrombolysis and recanalization of the graft. Pulmonary angiography via femoral access demonstrated complete occlusion of the right pulmonary artery (Figure 1b). Attempted access to the occluded right pulmonary artery graft for recanalization with multiple catheters and wires was unsuccessful. An ultrasound-assisted 6 French infusion catheter (EKOS, South Bothell, Washington) was placed through the main pulmonary artery, and tissue plasminogen activator (tPA) (Genentech, San Francisco, CA) was infused at 2 mg/h.

Repeat pulmonary artery angiogram through the existing infusion catheter 24 h later showed persistent occlusion of the right pulmonary artery graft and no significant change. Multiple catheters and wire combinations were used; however, the occluded right pulmonary artery graft could not be crossed. A 5 French Mikaelsson infusion catheter (Angiodynamics, Latham, New York) was used to infuse tPA at 1 mg/h (Genentech, San Francisco, CA). Heparin infusion was administered through both sheaths at 100 units/h.

On the following day, the 5 French Mikaelsson infusion catheter was exchanged for a 5 French pigtail catheter, and a pulmonary angiogram again showed absent flow into the main right pulmonary artery. Over a 0.035" guidewire (Cook Medical Inc., Bloomington, IN), a 6 French AR1 and guide catheter (Medtronic, Minneapolis, Minnesota) were used to engage a tiny crevice, created after the tPA chipped away at the occlusive thrombus, into the right pulmonary artery. Next, a 0.014" Confianza wire (Abbott, Abbott Park, Illinois) and 1.7 French microcatheter combination (Terumo, Somerset, New Jersey) were used to cross the occluded right pulmonary artery graft. An angiogram was performed, and the right pulmonary artery and branches were visualized. Several attempts to cross with a larger catheter were made, using both a 6 French EKOS infusion catheter and 4 French Fountain infusion catheter. Following the failure to cross with larger catheters, balloon angioplasty was performed using a 2 mm × 2 cm diameter balloon (Powercross-eV3/Medtronic, Minneapolis, MN), and subsequently, a 4 mm × 4 cm angioplasty balloon. Follow-up angiogram showed significant recoil following balloon dilation. A 4 French glide catheter was modified by poking holes in its most proximal 3 cm. Over a 0.035" extending glide wire, the modified 4 French glide catheter was buried at the anastomosis of the occluded right pulmonary artery graft, and tPA was infused a 1 mg/h.

An angiogram through the modified 4 French glide catheter was performed 24 h later and showed patent left pulmonary artery branches and residual thrombus in the right pulmonary graft. Stenosis was also noted through the graft. After numerous attempts, a 0.018 guidewire was advanced into the right lower lobe pulmonary artery branch. A 6 mm balloon

catheter was advanced through the graft and angioplasty was performed (Dorado - Bard Medical Division, Covington, GA) (Figure 2a). A 12 mm × 4 cm stent was deployed into the right pulmonary artery graft and dilated using the 6 mm × 4 cm balloon (Figure 2b). Subsequent angiography showed moderate improvement in arterial flow, although there was persistent stenosis immediately distal to the stent. Subsequent attempts to advance the balloon through the area of stenosis were unsuccessful. Catheter-directed thrombolysis (CDT) was continued per protocol with tPA infusion at 1 mg/h.

On day 4, following overnight catheter-directed thrombolysis, the pulmonary angiogram showed markedly improved flow to the distal branches with persistent focal stenosis at the distal




Figure 1: A 41-year-old female with anomalous aortic origin of the left coronary artery, status post-mobilization of the anomalous left main coronary artery, pulmonary artery translocation, and interposition of a 16 mm Dacron graft to the right pulmonary artery who presented with chest pain and shortness of breath. (a) Coronal image from a contrast-enhanced computed tomography of the chest demonstrating the main pulmonary artery and truncated the right pulmonary artery graft (circle). (b) Day 0: Pulmonary artery angiogram opacifying the native left pulmonary artery and truncated the right pulmonary artery graft (circle).




Figure 2: A 41-year-old female with anomalous aortic origin of the left coronary artery, status post-mobilization of the anomalous left main coronary artery, pulmonary artery translocation, and interposition of a 16 mm Dacron graft to the right pulmonary artery who presented with chest pain and shortness of breath. (a) Day 3: Right pulmonary artery angiogram demonstrating partially recanalized the right pulmonary artery branches (arrows). (b) Day 3: Right pulmonary artery angiogram of showing deployment of a 12 mm × 4 cm self-expanding, uncovered stent into the right pulmonary artery graft, and dilated using the 6 mm × 4 cm balloon (arrow).

end of the stent. The stenosis was further treated with an 8 mm × 4 cm balloon catheter (Dorado - Bard Medical Division, Covington, GA). Repeat angiography showed improved blood flow through the graft. The patient was subsequently discharged on aspirin, clopidogrel, and warfarin.

Follow-up was done 6 weeks later with CT angiography, which showed persistent stenosis of the right pulmonary artery graft distal to the stent. Angioplasty was again performed throughout the graft and a few centimeters distal where the stenosis was initially seen (Figure 3a). Follow-up angiogram performed 10 days later demonstrated markedly improved blood flow through the graft and pulmonary artery branches (Figure 3b).

The patient continues to be followed by the interventional radiology department with CT angiography and repeated angioplasty and stenting to treat the chronic pulmonary artery graft occlusion.

DISCUSSION

Coronary artery anomalies (CAAs) in one or more of the coronary arteries may be incidentally found in 0.3–1% of healthy individuals. Of the three types of CAAs, origin of left CAAs comprises 1.2% of CAA making the prevalence of AAOLCA 0.0036–0.012% in the general population.^[4] Management of AAOLCA is still debated, but definitive surgical treatment is recommended even for asymptomatic patients. Several different management options include coronary ostial reimplantation, unroofing of impinged coronary artery segment, and coronary




Figure 3: A 41-year-old female with anomalous aortic origin of the left coronary artery, status post-mobilization of the anomalous left main coronary artery, pulmonary artery translocation, and interposition of a 16 mm Dacron graft to the right pulmonary artery who presented with chest pain and shortness of breath. (a) 6-week follow-up right pulmonary artery angiogram after stent deployment and angioplasty with an 8 mm × 4 cm balloon catheter. There is improved contrast opacification of the right pulmonary artery branches and persistent residual stenosis (arrow) in the proximal right pulmonary artery stent graft. (b) 7-week follow-up right pulmonary artery angiogram after balloon angioplasty with 12 mm × 4 cm balloon demonstrating patent flow in the right pulmonary artery stent graft and pulmonary artery branches. Minimal residual stenosis was present at the end of the study (arrow)

stent deployment.^[5,6] However, due to the limited number of patients and debates of outcomes and limited information on long-term follow-up, large series of patients are needed to determine a standard of care.^[5]

During the surgical repair of any vessel, stenosis is a complication that is not unexpected. Our patient's RPA stenosis was treated with a Dacron stent. Dacron graft patency is well documented in peripheral bypass grafts; however, patency of pulmonary arterial grafts is not well elucidated. A contemporary meta-analysis of femoropopliteal bypass grafts found patency to be 60.2% at 3 years and 49.2% at 5 years.^[7] Furthermore, aortofemoral bypass grafts have shown to have 9.4% yearly thrombosis rate and a 5 year patency of 59%.^[8] Some of the various techniques for endograft recanalization are included as CDT, Fogarty thrombectomy, and laser recanalization.^[9,10] Thrombolysis has been documented in the literature as an effective means of recanalization. However, CDT often requires prolonged infusions with potential hemorrhagic complications.^[10,11] Fogarty thrombectomy has also been successfully used for endograft recanalization in peripheral endografts with a 71%–89% success rate. However, there was distal embolization in up to 9% of cases requiring adjunctive thrombolysis in 37% of patients.^[10-12] Laser recanalization of Dacron grafts has a limited success rate of 30% when compared to polytetrafluoroethylene grafts and has shown to have an increased residual stenosis average of 60%.^[13] Our recanalization technique using staged CDT and balloon angioplasty demonstrates a successful case for recanalization of occluded stent grafts in pulmonary arteries that could be used as a model for future cases.

CONCLUSION

While the technique for recanalization of peripheral arterial grafts is well documented, revascularization of pulmonary artery endografts is not well documented in the literature. In this case, the crossing was performed as done for any other CTO like that of peripheral arteries, in a staged approach. Our case illustrates a successful approach to the revascularization of an occluded pulmonary artery graft utilizing balloon angioplasty and CDT. Our case provides guidance of steps of crossing an occluded pulmonary artery graft to create reperfusion to pulmonary vasculature.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Yildiz A, Okcun B, Peker T, Arslan C, Olcay A, Vatan MB. Prevalence of coronary artery anomalies in 12,457 adult patients who underwent coronary angiography. *Clin Cardiol* 2010;33:E60-4.
2. Angelini P. Novel imaging of coronary artery anomalies to assess their prevalence, the causes of clinical symptoms, and the risk of sudden cardiac death. *Circ Cardiovasc Imaging* 2014;7:747-54.
3. Kalfa D. Postoperative pulmonary artery stenosis: Current options and future directions. *Transl Pediatr* 2017;6:57-8.
4. Zukić F, Miljko M, Vegar-Zubović S, Behmen A, Arapović AK. Prevalence of coronary artery anomalies detected by Coronary CT Angiography in Canton Sarajevo, Bosnia and Herzegovina. *Psychiatr Danub* 2017;29 Suppl 4:830-4.
5. Yuan SM. Anomalous origin of coronary artery: Taxonomy and clinical implication. *Rev Bras Cir Cardiovasc* 2014;29:622-9.
6. Elhmidi Y, Nöbauer C, Hörer J, Schreiber C, Lange R. Surgical unroofing of an anomalous right coronary artery arising from the posterior left sinus of valsalva. *World J Pediatr Congenit Heart Surg* 2013;4:433-5.
7. Takagi H, Goto SN, Matsui M, Manabe H, Umemoto T. A contemporary meta-analysis of Dacron versus polytetrafluoroethylene grafts for femoropopliteal by pass grafting. *J Vasc Surg* 2010;52:232-6.
8. Nevelsteen A, Suy R. Graft occlusion following aortofemoral Dacron bypass. *Ann Vasc Surg* 1991;5:32-7.
9. Pappy R, Hanna EB, Hennebry TA. Bifurcated aortic endograft limb occlusion managed with a novel method of isolated pharmacomechanical thrombectomy. *Catheter Cardiovasc Interv* 2010;76:895-900.
10. Amesur NB, Zajko AB, Makaroun MS. Treatment of a failed bifurcated abdominal aortic stent graft with thrombolysis and wallstent placement. *J Vasc Interv Radiol* 1997;8:795-8.
11. Silva JA, Ramee SR, Collins TJ, Jenkins JS, Lansky AJ, Ansel GM, *et al.* Rheolytic thrombectomy in the treatment of acute limb-threatening ischemia: Immediate results and six-month follow-up of the multicenter AngioJet registry. Possis Peripheral AngioJet Study AngioJet Investigators. *Cathet Cardiovasc Diagn* 1998;45:386-93.
12. Ansel GM, George BS, Botti CF, McNamara TO, Jenkins JS, Ramee SR, *et al.* Rheolytic thrombectomy in the management of limb ischemia: 30-day results from a multicenter registry. *J Endovasc Ther* 2002;9:395-402.
13. Kaelin LD, Klingman N, Normann SJ, Seeger JM. *In vitro* laser recanalization of chronically occluded prosthetic grafts. *Lasers Surg Med* 1992;12:410-6.

How to cite this article: Salamo R, Piening N, Garrity J, Moreno I, Vaheesan K. Revascularization of Chronic Occlusive Disease of a Right Pulmonary Artery Graft. *Am J Interv Radiol* 2018, 2(4) 1-4